

University of Groningen

Carbohydrates in the North Sea during spring blooms of *Phaeocystis*

Janse, Ingmar; van Rijssel, Marion; Gottschal, Jan C; Lancelot, C; Gieskes, Winfried W C

Published in:
Aquatic Microbial Ecology

DOI:
[10.3354/ame010097](https://doi.org/10.3354/ame010097)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1996

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Janse, I., van Rijssel, M., Gottschal, J. C., Lancelot, C., & Gieskes, W. W. C. (1996). Carbohydrates in the North Sea during spring blooms of *Phaeocystis*: A specific fingerprint. *Aquatic Microbial Ecology*, 10(1), 97-103. <https://doi.org/10.3354/ame010097>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Carbohydrates in the North Sea during spring blooms of *Phaeocystis*: a specific fingerprint

Ingmar Janse^{1,*}, Marion van Rijssel², Jan C. Gottschal¹, Christiane Lancelot³,
Winfried W. C. Gieskes²

Departments of ¹Microbiology and ²Marine Biology, University of Groningen, PO Box 14, 9750 AA Haren, The Netherlands

³Groupe de Microbiologie des Milieux Aquatiques, Université Libre de Bruxelles, Campus de la Plaine CP 221,
Boulevard de Triomphe, B-1060 Bruxelles, Belgium

ABSTRACT: Regional and temporal variation in the composition of water-soluble carbohydrates from *Phaeocystis* colonies sampled in the southern North Sea was small during spring 1994, except for a high variability in the contribution of glucose. Glucose is universally present in storage products of microalgae; the relative constancy of the carbohydrate pattern of the other monosaccharides suggests that these are part of the more refractory colony mucus. In all *Phaeocystis* samples arabinose dominated, followed by xylose (Belgian coast) or galactose and mannose (Dutch coast). Rhamnose, glucuronate and O-methylated sugars were present in lower amounts. The latter, always present in samples containing *Phaeocystis*, may be typical for North Sea strains. The sugar patterns we report here differ from those presented in the literature concerning *Phaeocystis*-derived material, and also from the sugar fingerprint in the preceding diatom bloom. The *Phaeocystis* mucus apparently behaves as particulate matter since it was retained on filters of over 1 µm. This characteristic together with its refractory nature, typical of 'transparent exopolymer particles' (TEPs), must have consequences for the heterotrophic microbial community in terms of adherence and substrate availability.

KEY WORDS: *Phaeocystis* · Mucus · Carbohydrates · Sugars

INTRODUCTION

The marine prymnesiophyte *Phaeocystis* has a world-wide distribution (Baumann et al. 1994). In Arctic and Antarctic regions, but also in the North Sea, it is capable of forming massive blooms that often dominate the plankton population (Gieskes & Kraay 1975, 1977, Lancelot et al. 1994). The taxonomic position and life cycle of *Phaeocystis* continue to be the subject of studies that started many decades ago (Kornmann 1955, Baumann et al. 1994, Medlin et al. 1994, Rousseau et al. 1994, Vaulot et al. 1994). The alga undergoes phase changes between flagellated single cells and non-motile cells organized in gelatinous colonies. The predominant form during the blooms is the colonial ('palmella') phase.

Colony mucus can contribute up to 90% to the total *Phaeocystis* biomass (Rousseau et al. 1990). This cell-surrounding gel is of a polysaccharide nature (Guillard & Hellebust 1971, Painter 1983). Guillard & Hellebust (1971) reported the carbohydrates galactose, glucose, mannose, arabinose, xylose, ribose, rhamnose and a hexuronic acid. The contribution of each sugar to the total amount was different in the Surinam and the Massachusetts (USA) strains that they analyzed. In a seawater sample dominated by *Phaeocystis*, Painter (1983) found a proteoglycan containing hemiester sulphate residues of galacturonic acid, glucosamine, galactose, arabinose, xylose, and rhamnose. The gelling properties of the matrix polysaccharides of *Phaeocystis* are promoted by salt bridges that involve calcium and magnesium ions that interlink the matrix polysaccharides which, by alcian blue staining, have been shown to contain carboxylated and sulphated groups (van Boekel 1992).

*E-mail: i.janse@biol.rug.nl

The physical properties of *Phaeocystis* colonies change through successive stages of the bloom, from regularly shaped colonies, essentially free of bacteria, to irregular colonies with sticky mucus (Verity et al. 1988, Rousseau et al. 1994, Thingstad & Billen 1994). Decaying colonies are often abandoned by the cells, leaving empty 'ghosts' that eventually turn into aggregates. It is likely that changes in the physical appearance of the colonies are reflected in their chemical structure and composition. The matrix composition and the possibly related firmness of colonies are of physiological interest since they may affect nutrient uptake and metal speciation (Lubbers et al. 1990), aggregation and sedimentation (Wassmann et al. 1990), and may even affect resistance to grazing by heterotrophs, ranging from protozoa to copepods (Weisse et al. 1994).

Heterotrophic bacteria interacting with the mucus might alter the matrix properties (Thingstad & Billen 1994). As a first step towards revealing possible *Phaeocystis*-bacteria interactions we investigated the carbohydrate composition of *Phaeocystis* colonies in the course of the spring bloom in the North Sea. Another purpose was to find a possible relationship between structural characteristics of the colony mucus matrix and its biochemical composition.

MATERIAL AND METHODS

Culturing conditions. *Phaeocystis* strain BCZ was isolated in 1993 by V. Rousseau at Stn 330 (51° 26' N, 2° 49' E; Fig. 1). The strain was cultured in 0.5 l serum bottles containing 0.4 l seawater ($S = 33\text{‰}$) from the

Dogger Bank area that was enriched with 'minor salts', trace elements, and nutrients according to Admiraal & Werner (1983) and also 0.4 ml of a vitamin solution (Veldhuis & Admiraal 1987) and additional bicarbonate ($0.2 \text{ g l}^{-1} \text{ NaHCO}_3$). Bottles were incubated in a culture cabinet (11°C, 14:10 L:D cycle) on a rolling device (8 rpm) at an irradiance of $40 \text{ mmol m}^{-2} \text{ s}^{-1}$, measured with a cosine collector.

Sampling. Samples were taken at regular intervals off the Belgian coast (Stn 330) and from the Marsdiep close to Texel, The Netherlands (Fig. 1). Samples were gently taken with a bucket in order to avoid colony disruption; they were then transferred to plastic bottles. In the laboratory, the samples were treated within 5 h in order to obtain the following fractions:

(1) Total organic matter, i.e. dissolved and particulate, obtained after homogenization of a frozen (-20°C) and thawed sample.

(2) Dissolved organic matter containing the $<1 \text{ }\mu\text{m}$ fraction obtained by applying gravity filtration, on preashed (4 h, 450°C) GF/C glass fibre filters (Whatman). Small volumes ($<10 \text{ ml}$) were filtered on relatively large filters (17.3 cm^3) to prevent clogging of the filters. With larger volumes the filters were changed regularly (vacuum pressure was never applied).

(3) Material in which colonies of *Phaeocystis* dominated was usually obtained by gentle filtration of 5 to 20 l over a $250 \text{ }\mu\text{m}$ mesh net. At some locations (Stns 800, dcz9, ro91, ro92, and pel94) material was obtained by towing a plankton net ($100 \text{ }\mu\text{m}$ mesh) through a dense *Phaeocystis* bloom ($>10^6 \text{ cells l}^{-1}$). Collected material was extracted for 5 min at 100°C after freezing and thawing. Samples were homogenized on a Vortex, centrifuged ($29\,000 \times g$ for 20 min), and dialyzed (Viskin size 1 8/32" Medicell International, pore size 1000 D) prior to sugar analysis.

For determination of the composition of polysaccharides present in the total seawater fraction, similar homogenization, centrifugation, and dialysis steps were performed prior to concentration using a tangential flow membrane filter (Mini-ultraset, Filtron Technology B.V., The Netherlands) with a pore size of 1 kD (4°C).

Phytoplankton abundance. The amount of phytoplankton cells was determined by counting of Lugol-preserved samples with an inverted microscope after gentle ultrasonic treatment of the samples to provide an even distribution of cells on the bottom of the counting vessel. Carbon biomass estimates were based on amount and size of cells and colonies. For *Phaeocystis* the calculation method described by Rousseau et al. (1990) was used, and for diatoms the factor of $0.11 \text{ pg C }\mu\text{m}^{-3}$ recommended by Edler (1979) was used.

Sugar analysis. Sugars present in a sample were analyzed by the modified phenol-sulphuric acid method described by Liu et al. (1973) with glucose as a refer-




Fig. 1. Sample sites along the North Sea coast

ence. For the determination of the relative sugar composition, samples were methanolized (2.0 M methanolic HCl, 24 h, 85°C) prior to analysis of the trimethylsilylated (N-reacylated) methyl glycosides on a Varian 3600 gas chromatograph equipped with a J&W DB-1 capillary column (diameter 0.32 mm, length 30 m) and a flame ionization detector. Identification of sugar derivatives was performed by comparison with known standards; it was confirmed by GLC/MS (Kamerling & Vliegenthart 1989). Unknown peaks were identified by GLC/MS as O-methylated hexoses and pentoses. These sugars were not available as reference sugars, so the molar ratio was estimated by assuming a degradation factor similar to those of hexoses and pentoses; this will result in an unknown systematic error in the exact amount of these methylated sugars.

RESULTS

At Stn 330 (Belgian coast) an unexpectedly dense diatom bloom (*Rhizosolenia delicatula* predominant)




Fig. 2. Stn 330 (Belgian coast), spring 1994. (A) Diatom and *Phaeocystis* colony biomass. Data from V. Rousseau. (B) Total amount of sugars measured in seawater and the <1 μm 'dissolved' fraction, expressed in glucose equivalents

was present in spring 1994 (Fig. 2A). Also near Texel the *Phaeocystis* bloom was preceded by this massive diatom bloom (Fig. 3). At both monitoring stations the phytoplankton community was dominated by *Phaeocystis* colonies only during a short period of time, from 28 April (Day 118) until late May. The unusually successful development of diatoms over *Phaeocystis* colonies in 1994 has been attributed by V. Rousseau (pers. comm.) to the very rainy weather conditions prevailing in the preceding winter, flushing more riverborne silicate into the North Sea than normal (Schaub & Gieskes 1991).

Based on microscopic observations, the *Phaeocystis* species blooming at all sites sampled in this study was identified as *Phaeocystis globosa* (Baumann et al. 1994). The variation in biomass and cell numbers of *Phaeocystis* at Stns 330 and tex are presented in Figs. 2A & 3A (data courtesy of V. Rousseau and G. C. Cadée). Maximum cell numbers at the Stns 330, tex, ro91, ro92, pel94, 800 and dcz9 were 6.5, 18.3, 52, 3.2, 1.0, 14.7 and $13.1 \times 10^6 \text{ cell l}^{-1}$, respectively.




Fig. 3. Stn tex (Texel, Dutch coast), spring 1994. (A) Diatoms and *Phaeocystis* colonial cells. Data from G. C. Cadée. (B) Total amount of sugars in the water column. All sugars were retained on GF/C filters




Fig. 4. Stn 330 (Belgian coast). (A) Upper panel: patterns of carbohydrates other than glucose measured in the $>250 \mu\text{m}$ fraction. Lower panel: percentage of glucose to total amount of sugars in the samples. (B) Upper panel: patterns of sugars other than glucose present in the total polysaccharide pool of the water column. Lower panel: percentage of glucose in the samples. Omp: O-methylated pentoses; ara: arabinose; rha: rhamnose; xyl: xylose; omh: O-methylated hexoses; man: mannose; gal: galactose; gla: glucuronic acid; fuc: fucose

High concentrations of total sugar were found during the spring phytoplankton growth phase, i.e. 0.7 to 7.5 mmol m^{-3} glucose equivalents (50 to 540 mg C m^{-3}) at the Belgian reference station (330). One peak coincided with the diatom bloom of *Rhizosolenia delicatula* (Day 111) and one with the *Phaeocystis* bloom (Days 131 and 138; Fig. 2A). At Stns 800 and dc9, 11.6 and 6.1 mmol m^{-3} were found, of which 100 and 89% , respectively, were in the fraction retained by GF/C filters. There was a notable shift from the sugars present predominantly in the 'dissolved' fraction, when the

diatom was blooming, to the 'particulate' fraction ($>1 \mu\text{m}$), i.e. total minus 'dissolved' when *Phaeocystis* became dominant (Fig. 2). Near Texel, all sugars were present in the $>1 \mu\text{m}$ fraction at concentrations varying between 1.9 and 14.9 mmol m^{-3} glucose equivalents (137 to 1070 mg C m^{-3} ; Fig. 3). The changes in the amounts of sugars in the water column corresponded to the changes observed in phytoplankton biomass (Fig. 3).

The *Phaeocystis* colony fractions that were obtained on a $250 \mu\text{m}$ mesh filter (or $100 \mu\text{m}$ in case of stations only sampled once) formed a green and viscous layer. The water-soluble carbohydrates that were extracted from this layer contained arabinose, rhamnose, xylose, mannose, galactose, glucose, and glucuronic acid as well as unidentified O-methylated pentoses and hexoses and traces of ribose (Figs. 4 to 6). N-acetyl-glucosamine was not found in these $>250 \mu\text{m}$ samples, indicating that the colony fractions were essentially free of chitin-containing plankton such as copepods.

The diatom bloom sugar pattern (fraction $>250 \mu\text{m}$; analyzed on Days 111 and 118 at Stn 330) differed considerably from that of the *Phaeocystis* bloom that followed. Only mannose, galactose and relatively large amounts of glucose (385 and 165 times the total amount of other sugars) were present; O-methylated sugars, always found during the *Phaeocystis* bloom, were lacking during the diatom bloom (Fig. 4A).

Glucose was also the dominant sugar in the total polycarbohydrate pool present in seawater (Fig. 4B), except for the period of the *Phaeocystis* bloom. At that time (Days 125 and 138) a number of sugars also found in the colony fraction (Fig. 4A) became predominant, which suggests that *Phaeocystis* contributed most to the polysaccharide pool.

The *Phaeocystis* bloom at Stn tex showed, apart from variable amounts of glucose, a rather constant pattern for the sugars in the $>250 \mu\text{m}$ colony fraction (Fig. 5). The glucuronic acid content seemed to increase slightly over time while mannose decreased. The anomalous pattern on Day 118 was possibly caused by aggregates of the waning *Rhizosolenia delicatula* bloom that were caught in the $>250 \mu\text{m}$ fraction.

The composition of non-glucose sugars in colony fractions of *Phaeocystis* sampled in 1994 at different locations were compared (Fig. 6). Arabinose was pre-

Fig. 5. Stn tex (Texel, Dutch coast). Upper panel: patterns of carbohydrates other than glucose measured in the >250 μ m fraction. Lower panel: percentage of glucose in the total amount of sugars in the samples. Abbreviations as in Fig. 4.





Fig. 6. Comparison of the carbohydrate patterns in different samples collected along the North Sea coast (stations as in Fig. 1). Glucose left out for reasons explained in the text. Abbreviations as in Fig. 3; lab: laboratory



dominant in all these fingerprints; the second most predominant sugar in the Belgian samples was xylose, of which the relative contribution to the total sugar amount decreased northward, while the relative amounts of galactose and mannose increased such that they became the second most predominant sugars in the Dutch samples. The O-methylated sugars were present in all *Phaeocystis*-containing samples, making up approximately 10% of the total amount of sugars (glucose not included). The samples ro91 and ro92, although taken in 1991 and 1992, resembled pel94 and tex94 taken in the same area. The laboratory culture of a strain isolated from Stn 330 (Belgian coast) in 1993 but cultured in water from the Dogger Bank (central North Sea) also resembled the Dutch fingerprints.

DISCUSSION

The sugar composition of the *Phaeocystis* colonies fraction revealed a complex fingerprint. Apart from the

highly variable amounts of glucose, the sugar pattern appeared to be similar along the entire continental North Sea coast, but it was completely different from those reported for *Phaeocystis* strains originating from the English Channel (Thingstad & Billen 1994), Surinam and Massachusetts (Guillard & Hellebust 1971), as well as from the diatom bloom preceding the *Phaeocystis* bloom (Fig. 4). Samples taken in the North Sea during the *Phaeocystis* bloom contained arabinose as the most abundant sugar, not mannose as reported by Guillard & Hellebust (1971). Furthermore, only traces of ribose and fucose were found in our samples. Instead, we identified O-methylated pentoses and hexoses in all our *Phaeocystis*-containing samples, including a laboratory culture originally isolated from Belgian coastal waters. These methylated sugars may be indicative of the *Phaeocystis* strains in the North Sea. The sugar pattern of the Belgian *Phaeocystis* samples is similar to that found by Haug et al. (1973) in a phytoplankton sample from the Trondheimsfjord on 30 March 1971, of which 63% of the cells was *Phaeocys-*

tis, except that the O-methylated sugars were not reported.

Despite the overall similarity of the fingerprints along the continental North Sea coast, some regional differences could be identified. In the Belgian samples, arabinose was twice as abundant as xylose and galactose, while in the Dutch area, mannose and galactose were most abundant, next to arabinose. Such local variability may reflect the presence of different strains. Cadée (1991) has suggested that regional blooms are inoculated by just a few colonies wintering in the area. This could be the mechanism by which local strains can develop. Strain differences in genome size and pigment contents have recently been reported for North Sea and English Channel isolates (Vaulot et al. 1994). Carbohydrate fingerprint differences might also be induced by environmental conditions (such as nutrient availability); an indication of this is the fingerprint of the strain isolated from the Belgian coast cultivated in water from the Dogger Bank area (central North Sea) which resembled the Dutch fingerprints.

The relative amounts of glucose were highly variable for the different sugar patterns, ranging from 0.2 to 380 times the amount of the other sugars. There are 3 possible explanations for this. Firstly, glucose polymers (glycogen, starch and laminaran) are universally produced as storage polymers by algae (Painter 1983). In fact, *Phaeocystis* produces a glucan as reserve polymer (Janse et al. 1996). The amount of glucose reserve polymers produced by algae depends highly on the environmental conditions (light and nutrients; Cuhel et al. 1984). Secondly, a contamination of the *Phaeocystis* colony fraction with other algae can cause a relatively high contribution of glucose to the sugar pool. Thirdly, the consumption of storage glucans can be expected to be rapid when they are consumed by bacteria that are adapted to this energy source. The amount of β -glucosidase at Stn 330 followed the total phytoplankton biomass development, not the seasonal pattern of any single species (S. Becquevort pers. comm.). Because processes that influence the glucose pool are not *Phaeocystis*-specific, glucose has been omitted from the comparison of sugar patterns here.

The relative amounts of sugars other than glucose present in *Phaeocystis* polymers were rather constant in the North Sea area. This suggests that these sugars are part of a structural rather than a storage polymer, needed to form the colony matrix. A structural polymer in *Phaeocystis* together with a storage glucan (Janse et al. 1996), resembles the situation in diatoms in which an acid-extractable storage glucan is commonly found in addition to more complex structural polymers (Haug & Myklestad 1976, Brockmann 1982, Hoagland et al. 1993).

The dramatic change of the colony shape during a bloom reported by Verity et al. (1988) and Rousseau et al. (1994) is not reflected in the sugar composition, which remained rather constant. Also, similar sugar fingerprints were found at the Stns 800 and dcz9, where healthy and decaying colonies dominated, respectively (V. Rousseau pers. comm.). Changes in colony firmness may depend on alterations in the amount of negatively charged carboxyl and sulphate groups (van Boekel 1992) or changes in polymer chain length and branching rather than on sugar composition per se.

The concentration of total sugars found during the *Phaeocystis* bloom is well within the range of 5 to 25 mmol m⁻³ glucose equivalents found earlier by Eberlein et al. (1985). Although *Phaeocystis* is known to be capable of high excretion rates of sugars (Guillard & Hellebust 1971, Lancelot & Mathot 1985, Veldhuis & Admiraal 1985) all carbohydrates were retained upon GF/C filters regardless of the state of the bloom that was sampled. In contrast, most of the carbohydrates present during the preceding diatom bloom passed these filters, except at the very beginning of the bloom when they were apparently associated with the cells. The reason for the different behavior of the carbohydrates produced by the diatoms and *Phaeocystis* probably lies in the production of matrix polysaccharides, which is so characteristic for *Phaeocystis*.

The constant sugar composition of this gel-forming colloidal material throughout the bloom may be indicative of its degradation-resistant nature, which results in a relatively long residence time. The result could be the formation of particles large enough to be retained on the filters, thus resembling the transparent exopolymer particles (TEPs) reported by Alldredge et al. (1993). The absence of dissolved sugars during and at the end of the *Phaeocystis* bloom can be explained by scavenging of dissolved organic matter by such particles (Decho 1990). Since bacteria have been shown to adhere to TEPs (Alldredge et al. 1993) and because the substrates available to the microbial loop are concentrated in these microenvironments (Azam & Smith 1991), the observation of a high concentration of TEP-like polysaccharides during *Phaeocystis* blooms in the North Sea suggests an important role of this mucus producing alga in the microbial loop.

Acknowledgements. We thank P. van der Gulik, V. Rousseau and R. Osinga for help in sampling, G. C. Cadée and V. Rousseau for biomass estimations, and Dr G. Gerwig (University of Utrecht, Bijvoet Centre) who performed the GC/MS measurements. Prof. J. P. Kamerling is acknowledged for access to the facilities in Utrecht. This work was supported by EC contract EV5V-CT94-0511, and by the Centre of Ecological and Evolutionary Studies (CEES) of the University of Groningen, The Netherlands.

LITERATURE CITED

- Admiraal W, Werner D (1983) Utilization of limiting concentrations of ortho-phosphate and production of extracellular organic phosphates in cultures of marine diatoms. *J Plankton Res* 5:495–513
- Aldredge AL, Passow U, Logan BE (1993) The abundance and significance of a class of large, transparent organic particles in the ocean. *Deep Sea Res* 40:1131–1140
- Azam F, Smith DC (1991) Bacterial influence on the variability in the ocean's biochemical state: a mechanistic view. In: Demers S (ed) *Particle analysis in oceanography*. NATO ASI Ser, Vol G27. Springer-Verlag, Berlin, p 213–236
- Baumann MEM, Lancelot C, Brandini FP, Sakshaug E, John DM (1994) The taxonomic identity of the cosmopolitan prymnesiophyte *Phaeocystis*; a morphological and eco-physiological approach. *J Mar Sys* 5:5–22
- Brockmann U (1982) Monosaccharide composition of acid-hydrolyzable carbohydrates in particulate matter during a plankton bloom. *Limnol Oceanogr* 27:770–776
- Cadée GC (1991) *Phaeocystis* colonies wintering in the water column? *Neth J Sea Res* 28:227–230
- Cuhel RL, Ortner PB, Lean DRS (1984) Night synthesis of protein by algae. *Limnol Oceanogr* 29:731–744
- Decho AW (1990) Microbial exopolymer secretions in ocean environments: their role(s) in food webs and marine processes. *Oceanogr Mar Biol A Rev* 28:73–153
- Eberlein K, Leal MT, Hammer KD, Hickel W (1985) Dissolved organic substances during a *Phaeocystis pouchetii* bloom in the German Bight (North Sea). *Mar Biol* 89:311–316
- Edler L (1979) Recommendations for marine biological studies in the Baltic sea. *Phytoplankton and chlorophyll*. Baltic Mar Biol Publ 5, Lund University, p 1–38
- Gieskes WWC, Kraay GW (1975) The phytoplankton spring bloom in Dutch coastal waters of the North Sea. *Neth J Sea Res* 9:166–196
- Gieskes WWC, Kraay GW (1977) Primary production and consumption of organic matter in the southern North Sea during the spring bloom of 1975. *Neth J Sea Res* 11:146–167
- Guillard RRL, Hellebust JA (1971) Growth and the production of extracellular substances by two strains of *Phaeocystis pouchetii*. *J Phycol* 7:330–338
- Haug A, Mykkestad S (1976) Polysaccharides of marine diatoms with special reference to *Chaetoceros* species. *Mar Biol* 34:217–222
- Haug A, Mykkestad S, Sakshaug E (1973) Studies on the phytoplankton ecology of the Trondheimsfjord. I. The chemical composition of phytoplankton populations. *J Exp Mar Biol Ecol* 11:15–26
- Hoagland KD, Rosowski JR, Gretz MR, Roemer SC (1993) Diatom extracellular polymeric substances: function, fine structure, chemistry, and physiology. *J Phycol* 29:537–566
- Janse I, van Rijssel M, van Hal PJ, Gerwig GJ, Gottschall JC, Prins RA (1996) The storage glucan of *Phaeocystis globosa* (Prymnesiophyceae) cells. *J Phycol* 32(3):in press
- Kamerling JP, Vliegthart JFG (1989) Mass spectrometry. In: Lawson AM (ed) *Clinical biochemistry — principles, methods, applications*, Vol 1, Mass spectrometry. Walter de Gruyter, Berlin, p 175–263
- Kornmann P (1955) Beobachtungen an *Phaeocystis*-Kulturen. *Helgoländer Wiss Meeresunters* 5:218–233
- Lancelot C, Mathot S (1985) Biochemical fractionation of primary production by phytoplankton in Belgian coastal waters during short- and long-term incubations with ^{14}C -bicarbonate. II. *Phaeocystis pouchetii* colonial population. *Mar Biol* 86:227–232
- Lancelot C, Wassmann P, Barth H (1994) Ecology of *Phaeocystis*-dominated ecosystems. *J Mar Sys* 5:1–4
- Liu D, Wong PTS, Dutka BJ (1973) Determination of carbohydrate in lake sediment by a modified phenol-sulfuric acid method. *Water Res* 7:741–746
- Lubbers GW, Gieskes WWC, del Castilho P, Salomons W, Bril J (1990) Manganese accumulation in the high pH microenvironment of *Phaeocystis* sp. (Haptophyceae) colonies from the North Sea. *Mar Ecol Prog Ser* 59:285–293
- Medlin LK, Lange M, Baumann MEM (1994) Genetic differentiation among three colony-forming species of *Phaeocystis*: further evidence for the phylogeny of the Prymnesiophyta. *Phycologia* 33:199–212
- Painter TJ (1983) Algal polysaccharides. In: Aspinall GO (ed) *The polysaccharides*, Vol 2. Academic Press, New York, p 96–285
- Rousseau V, Mathot S, Lancelot C (1990) Calculating carbon biomass of *Phaeocystis* sp. from microscopic observations. *Mar Biol* 107:305–314
- Rousseau V, Vaulot D, Casotti R, Cariou V, Lenz J, Gunkel J, Baumann MEM (1994) The life cycle of *Phaeocystis* (Prymnesiophyceae): evidence and hypotheses. *J Mar Sys* 5:23–29
- Schaub BEM, Gieskes WWC (1991) Eutrophication of the North Sea: the relation between Rhine river discharge and chlorophyll-a concentration in Dutch coastal waters. In: Elliot M, Ducrot JP (eds) *Estuaries and coasts: spatial and temporal intercomparisons*. Olsen and Olsen, Fredensborg, p 85–90
- Thingstad F, Billen G (1994) Microbial degradation of *Phaeocystis* material in the water column. *J Mar Sys* 5:55–65
- Van Boekel WHM (1992) *Phaeocystis* colony mucus components and the importance of calcium ions for colony stability. *Mar Ecol Prog Ser* 87:301–350
- Vaulot D, Birrien JB, Marie D, Casotti R, Veldhuis MJW, Kraay GW, Chrétiennot-Dinet MJ (1994) Morphology, ploidy, pigment composition, and genome size of cultured strains of *Phaeocystis* (Prymnesiophyceae). *J Phycol* 30:1022–1035
- Veldhuis MJW, Admiraal W (1985) Transfer of photosynthetic products in gelatinous colonies of *Phaeocystis pouchetii* (haptophyceae) and its effect on the measurement of excretion rate. *Mar Ecol Prog Ser* 26:301–304
- Veldhuis MJW, Admiraal W (1987) Influence of phosphate depletion on the growth and colony formation of *Phaeocystis pouchetii*. *Mar Biol* 95:47–54
- Verity PG, Villareal TA, Smayda TJ (1988) Ecological investigations of blooms of colonial *Phaeocystis pouchetii* — I. Abundance, biochemical composition, and metabolic rates. *J Plankton Res* 10:219–248
- Wassmann P, Vernet M, Mitchell BG, Rey F (1990) Mass sedimentation of *Phaeocystis pouchetii* in the Barents Sea. *Mar Ecol Prog Ser* 66:183–195
- Weisse T, Tande K, Verity P, Hansen F, Gieskes W (1994) The trophic significance of *Phaeocystis* blooms. *J Mar Sys* 5:67–79